arXiv:math/0305196v1 [math.MG] 14 May 2003

Infinite serie of extreme Delaunay polytope

Mathieu DUTOUR
LIGA, ENS/CNRS, Paris and
Hebrew University, Jerusalem*

February 1, 2008

Abstract

A Delaunay polytope P is said to be *extreme* if the only (up to isometries) affine bijective transformations f of $\mathbb{R}^n$, for which $f(P)$ is again a Delaunay polytope, are the homotheties. This notion was introduced in [DGL92]; also some examples in dimension 1, 6, 7, 15, 16, 22, 23 were constructed and it was proved that in dimension less than 6 there are no extreme Delaunay polytopes, except the segment. In this note, for every $n \geq 6$ we build an extreme Delaunay polytope ED_n of dimension n.

Let $L \subset \mathbb{R}^n$ be a n-dimensional lattice and let $S = S(c, r)$ be a sphere in $\mathbb{R}^n$ with center c and radius r. Then, S is said to be an *empty sphere* in L if the following two conditions hold:

$$\|v - c\| \geq r \text{ for all } v \in L \text{ and the set } S \cap L \text{ has affine rank } n+1.$$

The polytope P, which is defined as the convex hull of the set $P = S \cap L$, is called a *Delaunay polytope*.

Denote by D_n the root lattice defined as

$$D_n = \{(x_1, \dots, x_n) \in \mathbb{Z}^n \text{ with } \sum_{i=1}^{n} x_i \text{ even}\} .$$

The Delaunay polytopes of this lattice are:

- *Half-cube* $\frac{1}{2}H_n = \{x \in \{0,1\}^n \text{ with } \sum_{i=1}^n x_i \text{ even}\}$; it has 2^{n-1} vertices.
- *Cross-polytope* $C_n = \{e_1 \pm e_i \text{ with } 1 \leq i \leq n\}$ with $e_i = (0, \dots, 1, \dots, 0)$; it has $2n$ vertices.

Define the polytope P_n to be the convex hull of the three following layers of points:

---

*Research financed by EC's IHRP Programme, within the Research Training Network "Algebraic Combinator ics in Europe," grant HPRN-CT-2001-00272.

- a vertex $V = (\frac{1}{2}, \dots, \frac{1}{2}, 1)$,
- 2^{n-2} vertices $(x_1, \dots, x_{n-1}, 0)$ with $x_i \in \{0, 1\}$ and $\sum_{i=1}^{n-1} x_i$ even,
- $2(n-1)$ vertices $V_{j,\pm} = (\frac{1}{2}, \dots, \frac{1}{2}, -1) \pm e_j$ with $1 \leq j \leq n-1$.

It is easy to see, that, denoting by L_n the lattice generated by all $(x, 0)$ with $x \in D_{n-1}$ and $(\frac{1}{2}, \dots, \frac{1}{2}, 1)$, the third layer belongs to L_n if and only if n is even.

Theorem 1 *Let n be even and greater than 6.*

(i) P_n is an extreme Delaunay polytope; the corresponding quadratic form is $q(x) = x_1^2 + \dots + x_{n-1}^2 + \frac{n-3}{4} x_n^2$.

(ii) The center of the circumscribing sphere is $(0, \dots, 0, \frac{-1}{n-3})$, it radius is $\frac{n-2}{\sqrt{n-3}}$.

(iii) P_6=Schläfli polytope; its group of isometries has size 51840 and is transitive on vertices. If $n > 6$, then its group of isometries is $Sym(n-1) \times 2^{n-2}$ and there are three orbits of vertices.

Proof. In order to prove extremality, we will use geometry of numbers, i.e. in order to show that the only transformations preserving the property of being Delaunay are modulo isometries and translations the homotheties, we will determine the set of quadratic forms q such that P_n is a Delaunay polytope for q.

The restriction of the quadratic form q to the plane $x_n = 0$ left us with the 2^{n-2} vertices $(x_1, \dots, x_{n-1}, 0)$ with $x_i \in \{0, 1\}$ and $\sum_i x_i$ even. The rank of Half-cube is n (see [DeLa97] and [Du]), we can write q restricted to the plane $x_n = 0$ as $\sum_{i=1}^{n-1} \alpha_i x_i^2$ with $\alpha_i > 0$. We write the form q in the following way:

$$q(x) = \sum_{i=1}^{n-1} \alpha_i (x_i + \beta_i x_n)^2 + \gamma x_n^2 \ .$$

Denote by $S(c, r)$ the empty sphere of center c and radius r circumscribing $\frac{1}{2} H_{n-1}$ and V. Denoting $c = (h_1, \dots, h_{n-1}, h_n)$, one obtains the equations:

$$\begin{cases} \sum_{i=1}^{n-1} \alpha_i (h_i - x_i + \beta_i h_n)^2 + \gamma h_n^2 = r^2, \\ \sum_{i=1}^{n-1} \alpha_i (h_i - \frac{1}{2} + \beta_i (h_n - 1))^2 + \gamma (h_n - 1)^2 = r^2 \ . \end{cases}$$

The first equation is satisfied for all vectors x_i with $\sum_{i=1}^{n-1} x_i$ even and $x_i \in \{0, 1\}$. This implies, since $n \geq 6$ the relation $h_i + \beta_i h_n = \frac{1}{2}$. So, the equations reduce to:

$$\begin{cases} \frac{1}{4} \sum_{i=1}^{n-1} \alpha_i + \gamma h_n^2 = r^2, \\ \frac{n-2}{2} \sum_{i=1}^{n-1} \alpha_i \beta_i^2 + \gamma (h_n - 1)^2 = r^2 \ . \end{cases}$$

Consider now the $2(n-1)$ vertices $V_{j,\pm} = (\frac{1}{2}, \dots, \frac{1}{2}, -1) \pm e_j$ with $1 \leq j \leq n-1$.

Since $V_{j,+}$, and $V_{j,-}$ are on the sphere $S(c, r)$, one obtains:

$$\begin{cases} \alpha_j (1 - \beta_j)^2 + \sum_{i \neq j} \alpha_i \beta_i^2 + \gamma (h_n + 1)^2 = r^2, \\ \alpha_j (1 + \beta_j)^2 + \sum_{i \neq j} \alpha_i \beta_i^2 + \gamma (h_n + 1)^2 = r^2 \ . \end{cases}$$

Above two equations yield $\beta_j = 0$. If all $V_{j,\pm}$ are on the sphere, then $\alpha_j = \alpha$ is independent of j and all equations simplify to:

$$\alpha + \gamma(h_n+1)^2 = r^2\ ,\ \ \frac{n-1}{4}\alpha + \gamma h_n^2 = r^2 \ \text{ and }\ \gamma(h_n-1)^2 = r^2\ ,$$

whose solution is:

$$h_n = \frac{1}{3-n},\ \ \gamma = \alpha\frac{n-3}{4},\ \ r = \frac{n-2}{\sqrt{n-3}}\sqrt{\alpha}\ \ .$$

So, q is equal to $\alpha(\sum_{i=1}^{n-1} x_i^2 + \frac{n-3}{4}x_n^2)$.

Now, let us prove that this is a Delaunay polytope. If v is an interior point of P_n, then it belongs to one of the three layers $x_n = -1,\ 0,\ 1$. But each one of the corresponding sections (i.e. Cross-polytope, Half-cube or point) is a Delaunay polytope; so, there is no such v.

Let us find the symmetry group. Pairwise distance of adjacent vertices of Half-cube is $\sqrt{2}$. Distance between V and vertices of Half-cube is $\sqrt{\frac{n-2}{2}}$. So, if $n > 6$, then any isometry preserving P_n must leave the vertex V invariant; hence, if $n > 6$, then the symmetry group is that of Half-cube, i.e. $Sym(n-1) \times 2^{n-2}$.

Standart computation shows that P_6 is the Schläfli polytope, whose symmetry group has size 51840. □

Define ED_n to be P_n if n is even and the polytope obtained from P_{n-1} by using Lemma 15.3.7 is f is odd. Remark that ED_6 (respectively ED_7) is Schläfli (respectively, Gosset) polytope.

Corollary 1 *If $n \geq 6$, then the polytope ED_n is extreme.*

Proof. If n is even, this follows from 1, while if n is odd, this follows from Lemma 15.3.7. □

Remark, that ED_6 is *unique* extreme Delaunay polytope in dimension 6 (see [DeDu]); *all* 6241 6-dimensional Delaunay polytopes were listed in [Du].

References

[DeLa97] M.Deza and M.Laurent, *Geometry of cuts and metrics*, Springer–Verlag, Berlin 1997.

[Du] M.Dutour, *The six-dimensional Delaunay polytopes*, submitted, `http://www.arxiv.org/abs/math.MG/0212353`.

[DeDu] M.Deza and M.Dutour, *The hypermetric cone on seven vertices*, submitted, `http://www.arxiv.org/abs/math.MG/0108177`.

[DGL92] M. Deza, V.P. Grishukhin, and M. Laurent, *Extreme hypermetrics and L-polytopes*, in G.Halász et al. eds *Sets, Graphs and Numbers, Budapest (Hungary), 1991*, **60** *Colloquia Mathematica Societatis János Bolyai*, (1992) 157–209.